

Research article

EFFECT OF TEMPERATURE AND RELATIVE HUMIDITY ON POPULATION DYNAMICS OF PREDATORS OF COTTON PESTS

*Muhammad Khan, Dr. Roshankhanzada, Tauqir Ahmad

*Corresponding Author= muhammadkhan1979@gmail.com



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

ABSTRACT

The study on effect of biotic factors on population dynamics of predators of cotton pests was conducted in farmer field school Bhattianwala, Punjab Pakistan during cotton growing season 2013. The population of predators was noted on weekly basis and data of temperature and relative humidity was taken from meteorological department. Data revealed that major predators of cotton was spider, ants, *chrysopia*, *campiloma* and orius bug. It was observed that there was positive correlation between ants and temperature (0.296), while there was negative correlation between temperature and *campiloma* (-0.699), spider and temperature (-0.4575) and there was also negative correlation between temperature and orius bug (0.11305). There was positive correlation between *Chrysopa* and temperature (0.543866). Humidity had positive correlation with orius bug (0.43058), spider (0.700366) and *campiloma* (0.835869) while it had negative correlation with ants (-0.21196), *chrysopa* (-0.37335). **Copyright © WJAFST, all rights reserved.**

INTRODUCTION

Cotton is a major cash crop in Pakistan contributing about 2 percent of GDP and 8.2 % of value added in Agriculture, with total area under crop 2.9 million hectare and production of 10.0 million bales average 571 kg seed cotton per hectare. In 1999-2000 the cotton production was 11.24 million bales which declines to 10.73 million bales in 2000-01, it further dipped to 10.61 million bales in fiscal year 2001-02 and 10.21 million bales in 2002-03, the downslide continued till 2003-04 reaching 10.0 million bales with 1.6% lower yield than the last year (Eco. Survey of Pak. 2003-04). The production of cotton in 2006-07 was 12.9 million bales which was decreased by 11.7 million bales in 2007-08. In 2010-11 the cotton production was estimated 11.46 million bales and the production of cotton crop has increased in 2011-12 valued as 13.59 million bales (Eco. Survey of Pak. 2011-12). Heavy insect/pest attack and lack of resistant varieties were the main causes of this decline. The country at present imports over 10 billion of pesticides annually through registered firms alone. About 90% of these are being used on cotton (Ingram, *et al.* 1989).

Thus there is a need to develop alternative methods. Biological control seems to have great potential in terms of profitability, safety for humans and animals and for environment and for the sustainability of agriculture activities. It requires the knowledge of biology, ecology, phenology and behavior of pests and their natural enemies (Mallah, *et al.* 2001). Biological control is the most important component of IPM because a number of pests of a crop, remain under natural control. In sampling southwestern desert cotton, Haggier and Narnjo, 1993 have identified approximately 25 species of predatory arthropods associated with spotted bollworms population. Field assessment revealed highest population of predators early in the season with rapid decline associated with pesticide application. *Chrysoperla carnea* and *Geocoris punctipes* were notable exceptions and were found to increase through the season. Among the most common predators were *Chrysoperla carnea*, *Groceries spp.*, *Oriustriticolor* and *Nabis alternatus*.

Mullah *et al.* 2001 described that the characteristics of predators that it should be able to survive in all the niches and throughout the climatic range occupied by the pest. The relationship of its development and voracity to temperature determines whether it can cause mortality sufficiently early in the pest annual cycle and whether it can avoid being outstripped by the pest. Keeping in view the importance of above facts and economic importance of Integrated Pest Management in the cotton crop, the study on appearance and abundance of various insect predators

(natural enemies) with relation to temperature and relative humidity was conducted during cotton growing season 2013.

MATERIALS AND METHODS

The trail was laid out at plot in Bhattianwala, a surrounding village of Tehsil Kabirwala District, Khanewal, Pakistan during the cotton growing season 2013. The trail was laid in a randomized complete block designed with three replicates. 1st CESA (Cotton Ecosystem Analysis) was taken on 19-06-2013 and last CESA was taken on 25-09-2013. Cotton ecosystem analysis comprised of weekly agronomical and entomological data was taken from selected twenty five plants from the whole plot. Agro meteorological data for temperature and relative humidity was taken from Agro metrological center Multan. Each plant was observed thoroughly, data for predators was collected from whole plant. Total number of predators were divided with total number of plants examined to get average per plant. At the end of season, the data was subjected for statistical analysis to get simple correlation between predators of insect pests of cotton and weather factors (temperature and relative humidity).

RESULTS AND DISCUSSION

Population dynamics of *Chrysopa* spp.

The population of *Chrysopa* (*Chrysoperacarnea*) was at its peak value 8.2 at 30 C⁰ and 77.45 relative humidity on last week of August highest temperature and lowest humidity. Data in table. 1 shows that population of *Chrysopa* is negatively correlated (-0.3733) with temperature and it is positively correlated with relative humidity (0.543886). The present findings are in agreement with those of Mari *et al.* (2006) who described that consolidated in population of predators was maximum on 13th September and showed there was direct relation of pest with population of predators. Radhakrishnan and Muraleedhnan (1994) observed the seasonal abundance of *C.carnea*. The same results were reported by Holz(1994). Similarly Atakan and Ogur (1994) studied the effect of *C.carnea*.

Population dynamics of *Campiloma*

The population of *Cmpiloma* was at its peak value 6.8 at 29.55 C⁰ and 81.6% relative humidity on last week of August (fig. 1) Table. 1 showed that temperature had negative correlation with population of *compiloma* (-0.699) and positive correlation with humidity (0.835869). Present finding are in agreement with those of Mari *et al.* (2006)

who described that consolidated in population of predators was maximum on 13th September and showed there was direct relation of pest with population of predators.

Population dynamics of Orius Bug

Population of Orius bug was at its peak value 10.1 at 32.5 C⁰ and 77.9% relative humidity on 2nd week of August (fig. 1) Table.1 showed that population of Orius bug has negative correlation with temperature (-0.11305) and positive correlation with humidity (0.43058). Present findings are in agreement with those of J.M. Mari *et al.* (2006) who described that consolidated in population of predators was maximum on 13th September and showed there was direct relation of pest with population of predators.

Population dynamics of Spider

Population of Spider was at its peak value 3.48 at 30.6 C⁰ and 77.4 5% relative humidity on 3rd week of August (fig. 1) Table. 1 showed that population of spider had negative correlation with temperature (-0.4575) and had positive correlation with humidity (0.700366). Present study was in conformation with Mao (1984) who reported that spiders were present from late June to late August.

Population dynamics of Ants

Population of ants was at its peak value 5.89 at 29.55 C⁰ and 81.6% relative humidity on 3rd week of August (fig. 1) Table. 1 showed that population of ants had positive correlation with temperature (0.296) and had negative correlation with humidity (-0.21196) present findings are in agreement with those of Mari,*et al.* (2006) who described that consolidated in population of predators was maximum on 13th September and showed there was direct relation of pest with population of predators.

Table 1. Correlation of Campiloma, Ants, Orius Bug, Spider and Chrysopasppa with Temperature and Humidity during cotton growing season 2006

| | Campiloma | Ants | Orius | Spider | Chrysopa |
|-------------|-----------|----------|----------|----------|----------|
| Humidity | 0.835869 | -0.21196 | 0.43058 | 0.700366 | -0.37335 |
| Temperature | -0.699 | 0.296 | -0.11305 | -0.4575 | 0.543886 |

Correlation at 5% level

LITERATURE CITED

Atakan, E. and Ogur, A.F. 1994. Effect of natural enemies on the population development of cotton aphid (*Aphis gossypii*Glov.) (Homoptera: Aphididae). *EgeUniversitesiFakultesi, BitkiKorumaBolumu, Izmir*, 459-470.

Eco. Survey of Pak. (2003-04). Eco. Advisor's wing Isd: 12-13 Fryxell P.A., Staten G., Proter J.H1958. Performance of some wide crosses in Gossipier. *New Max AgriSta Bull* 4191-15.

Highlights of Eco. Survey of Pak. (2007-08).

Highlights of Eco. Survey of Pak. (2011-12).

Hagler, J. R. and S.E. Naranjo. 1993. An over view of biological control of sweet potato white fly *Bemesiatabaci*. USDA-ARS cotton Res. Lab. 4135. Broadway, Phoenix, AZ. Proceeding Beltwide cotton conferences, 2:680-685.

Mallah, G.H., A.K.Korejo, A.R. Soomro and A.W. Soomro. 2001. Population dynamics of predatory insects and biological control of cotton pest in Pakistan. *Online J. Biol. Sci.* 1 (4):245-248.

Holz, 1994. On the occurrence of stenophagous predators of aphids in winter wheat fields nearby Halle (Saxony-ANHALT). *Bull OILB SROP.* 17(4)127-136.

Ingram, W.R., J. A. Sutherland, H.Haque and I. D.J.De.B.1989. Pesticide Management in cotton in Pakistan. Ingree, M.Band Lyon, D.J. De.B (Eds.) *Pest Management in cotton* Elis Horwood, Chichester, U.K.

Mari, J.M., S. M. Nizamani and M.K. Lohar 2006. Population fluctuation of sucking insect pests and predators in cotton ecosystem. *A.C.S.C. proceeding* (8):929-934.

Mao, G.H and Z.C.Xia, 1984. Observation on the population dynamics of natural enemies of the cotton aphids on cotton. *Insects Knowledge (KunchongZhishi)* 1983. 20:217-219. Junshan State farm, Hunan, China. *RAE.* 1984. 72(12), 8043.

Radhakrishana, B. and Muraleedhran, N. 1994. Bio-ecology of six species of syrphid predators of tea aphid, *Toxopeteraaurantii* (Boyer de Fonscolombe) in Southern Ind., *Entomology.* 18(3):175-180.